arXiv:1211.5392v1 [math.AP] 22 Nov 2012

An approximate treatment of gravitational collapse

Yago Ascasibar[1], Rafael Granero-Belinchón[2] and José Manuel Moreno

November 26, 2012

Abstract

This work studies a simplified model of the gravitational collapse of a protostar based on the assumption that the mean velocity of the gas is proportional to the local acceleration. This assumption leads to a nonlocal generalization of the restricted Patlak-Keller-Segel model. We discuss some of its fundamental properties and illustrate them by means of numerical simulations. In particular, we are interested in the conditions for the gravitational stability of an infinite gas cloud, represented by $\mathbb{T}^d$. Local well-posedness in Sobolev spaces is proven for non-local pressure forces. Moreover, we show the smoothing effect of our equation, as well as a *Beale-Kato-Majda*-type criterion in terms of $\|\rho\|_{L^\infty}$. We also show that the problem is ill-posed in Sobolev spaces when it is considered backward in time. Finally, we prove that, in the critical case (one conservative and one dissipative derivative), $\|\rho\|_{L^\infty}(t)$ is uniformly bounded in terms of the initial data for sufficiently large pressure forces.

Keywords: Gravitational collapse, Star formation, Patlak-Keller-Segel model, Fractional calculus, Well-posedness, Instant analyticity, Blow-up, Simulation.

Acknowledgments: Y. Ascasibar receives financial support from project AYA 2010-21887-C04-03 and the *Ramón y Cajal* programme of the Spanish *Ministerio de Ciencia e Innovación* (MICINN). R. Granero-Belinchón is supported by grant MTM2008-03754 of the MICINN.

Contents

1 Introduction **2**

2 Well-posedness **5**
  2.1 Well-posedness in Sobolev spaces 5
  2.2 Smoothing effect 9

3 Gravitational stability **13**
  3.1 Minimum time for star formation 13
  3.2 Exponential suppresion of density fluctuations 14

---

[1] Email: `yago.ascasibar@uam.es`
Universidad Autónoma de Madrid
Departamento de Física Teórica
Campus de Cantoblanco, 28049 - Madrid (Spain)

[2] Email: `r.granero@icmat.es`
Consejo Superior de Investigaciones Científicas
Instituto de Ciencias Matemáticas (CSIC-UAM-UC3M-UCM)
C/Nicolás Cabrera, 13-15, Campus de Cantoblanco, 28049 - Madrid (Spain)

4 Numerical simulations **17**

1 Introduction

Gravitational collapse - the infall of a body under its own gravity - is of paramount importance in Astrophysics. Stars and galaxies grow from initially small preturbations of a uniform density medium, which is an unstable equilibrium solution of the Newtonian equations of motion. Upon the action of gravity, overdense regions attract the neighbouring material, accreting all mass within a certain radius of influence into a singular point. Some of the gravitational energy, though, transforms into heat and/or random motions that oppose collapse. In most real-life situations (black holes being the exception), the ensuing pressure gradient is able to counteract the gravitational acceleration, and a non-singular equilibrium state is reached.

Mathematically, the evolution of a fluid with mass density $\rho(\mathbf{r},t)$ and velocity $\mathbf{v}(\mathbf{r},t)$ is described by the Euler equations, that amount to the conservation of mass

$$\partial_t \rho + \nabla \cdot (\rho \mathbf{v}) = 0 \tag{1}$$

and momentum

$$\partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla)\, \mathbf{v} = -\frac{\nabla P}{\rho} - \nabla U \tag{2}$$

supplemented by the Poisson equation

$$\Delta U = 4\pi G\, (\rho - \langle \rho \rangle) \tag{3}$$

that relates the gravitational potential $U(\mathbf{r},t)$ to the density distribution of the fluid, as well as by an equation of state that specifies the function $P(\rho)$ or, equivalently, the sound speed

$$c_{\mathrm{s}}^2 = \frac{\partial P}{\partial \rho} \tag{4}$$

where P denotes the gas pressure. Finally, the initial data in ρ and $\mathbf{v}$ at some arbitrary time t_0 have to be specified in order to close the system.

The term $\langle \rho \rangle$ in equation (3) corresponds to the average gas density, and it vanishes for any system with finite mass contained in an infinite space. Here, we will consider the collapse of a protostar out of a gas cloud that is assumed to be roughly uniform on scales much larger than those relevant to the problem. Such conditions may be realized by a periodic cubic box of sufficiently large size L, i.e. a three-dimensional torus $\mathbb{T}^3$, as the spatial domain. In general, for a d-dimensional torus with periodicity L,

$$\langle \rho \rangle = \frac{1}{L^d} \int_{\mathbb{T}^d} \rho(x)\ dx. \tag{5}$$

We will also make the approximation, akin to Darcy's Law in Earth sciences, that the acceleration term that appears in the momentum conservation equation is always proportional to the fluid velocity

$$\partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla)\, \mathbf{v} \approx \frac{\mathbf{v}}{\tau} \tag{6}$$

where τ is a constant with dimensions of time. Substituting in expression (2) and taking the divergence, one obtains

$$\frac{1}{\tau} \nabla \cdot (\rho \mathbf{v}) = -\Delta P - \nabla\, (\rho \nabla U)\,,$$

which, using the continuity and Poisson equations and writing the equation of state in terms of the sound speed, transforms into

$$\frac{\partial_t \rho}{\tau} = c_{\mathrm{s}}^2 \Delta\rho + \nabla\rho\,\nabla U + 4\pi G\rho\left(\rho - \langle\rho\rangle\right).$$

Since U is fully determined by ρ, our model consists on a single, non-linear and non-local, partial differential equation that describes the evolution of the gas density.

In order to set the constant τ, let us consider the collapse of a spherically-symmetric perturbation where the density decreases monotonically with radius. This will always be the case in the vicinity of a density peak, and, for Gaussian random fluctuations, it will also hold, on average, at large radii [1]. Under those conditions, the different spherical shells will not cross during the collapse phase, and the enclosed mass $M(r)$ will be conserved for every shell. Starting from an initially homogeneous configuration,

$$\frac{|\mathbf{v}|}{|\partial_t \mathbf{v} + (\mathbf{v}\cdot\nabla)\,\mathbf{v}|} = \frac{\sqrt{\frac{2GM}{r}}}{\frac{GM}{r^2}} = \sqrt{\frac{2r^3}{GM}} = \sqrt{\frac{3}{2\pi G\rho}},$$

so we adopt

$$\tau = \sqrt{\frac{3}{2\pi G\langle\rho\rangle}} \tag{7}$$

in order to recover the exact solution at arbitrarily small times.

Denoting the solution of $\Delta u = f - \langle f\rangle$ by $u = T(f)$, defining $\beta = \frac{\pi c_{\mathrm{s}}^2}{G\langle\rho\rangle L^2}$, and choosing the units of time, length, and mass such that $\frac{1}{\sqrt{24\pi G\langle\rho\rangle}} = 1$, $L = 2\pi$, and $\langle\rho\rangle = 1$, respectively, we finally arrive at

$$\partial_t\rho = \beta\Delta\rho + \rho(\rho-1) + \nabla\rho\cdot\nabla T(\rho). \tag{8}$$

The case $\beta = 0$ would be a model for a perfectly collisionless fluid (e.g. cold dark matter), whereas $\beta > 0$ corresponds to an isothermal ideal gas. Moreover, we consider the generalized equation

$$\partial_t\rho = -\beta\Lambda^\alpha\rho + \rho(\rho-1) + \nabla\rho\cdot\nabla T(\rho) \tag{9}$$

where the fractional Laplacian $\Lambda^\alpha = (-\Delta)^{\alpha/2}$ is defined using Fourier theory:

$$\widehat{\Lambda^\alpha u} = |\xi|^\alpha \hat{u}.$$

Phisically, this is a nonlocal adhesion model. Due to the nonlocal character of the diffusive operators Λ^α, the pressure forces now contain information about the total distribution of mass; in other words, the pressure at a given point x does not depend only on $\rho(x)$ but on the overall distribution $\rho(y)$ for all y. Such an equation may develop singularities in a finite time depending on the initial data. The case $\alpha = 1$ is critical in the sense that we have a conservative derivative (the term $\nabla\rho\cdot\nabla U$) and a *'dissipative'* one in the term $-\beta\Lambda\rho$.

Equation (8) can be re-written as

$$\partial_t\rho = \beta\Delta\rho + \nabla\cdot(\nabla U\rho), \quad \Delta U = \rho - \langle\rho\rangle,$$

that bears obvious resemblances to the vorticity equation for the incompressible Navier-Stokes equations in two dimensions

$$\partial_t\omega = \nu\Delta\omega + \nabla\cdot(\nabla^\perp\psi\omega), \quad \Delta\psi = \omega,$$

and it has been proposed by Jäger and Luckhaus [20] as a simplified version of the classical (parabolic-elliptic) Patlak-Keller-Segel model [27, 22] of chemotaxis in biological systems (see e.g. [8, 10, 11, 12] and references therein; a recent review of the results concerning this equation can be found in [9]). The non-periodic (bounded or otherwise) case with $\langle\rho\rangle = 0$ has been previously studied in the context of gravitational collapse by P. Biler and co-workers, who have considered the problems of existence, conditions for blow-up, radial solutions, stationary solutions, and other qualitative properties [4, 5, 6]. L. Corrias, B. Perthame and H. Zaag have proved that, for small data in $\|\rho_0\|_{L^{d/2}}$, there are global in time weak solution to equation (8) in $d > 1$, and blow up if the smallness condition does not hold (see Theorem 1.1 and 1.2 in [18]). The periodic case has been studied by T.Senba and T.Suzuki in [29]. In this paper they consider the case of measure valued weak solutions.

We argue that there are several reasons why this equation constitutes a valid approach to the problem of gravitational collapse seeded by small perturbations of an initially homogeneous medium in equilibrium (a situtation often encountered in Astrophysics). First, the family of hydrostatic equilibrium solutions of the full problem,

$$\nabla P + \rho \nabla U = 0,$$

are also equilibrium solutions of our approximation, since, taking the divergence of the above expression,

$$\beta \Delta \rho + \rho(\rho - 1) + \nabla\rho \cdot \nabla T(\rho) = 0.$$

In addition, the model is also consistent with more classical alternatives based on linear perturbation theory, being, in fact, very similar in spirit (though not exactly equivalent) to the Zel'dovich approximation [30] used in Cosmology to study the formation of large-scale structure in the primordial universe. In this approximation, particles move along straight trajectories of the form

$$\mathbf{r}(t) = a(t)\,\mathbf{q} + b(t)\,\mathbf{p}(\mathbf{q})$$

where $\mathbf{r}$ denotes the actual (Eulerian) position of the particle, $\mathbf{q}$ is its Lagrangian coordinate, the factor $a(t)$ accounts for cosmic expansion, and the vector function $\mathbf{p}(\mathbf{q})$ is determined by the initial conditions. The evolution of the inhomogeneities is given by the growth factor $b(t)$, chosen to match the results of Eulerian linear theory. For an Einstein-deSitter cosmology, which always provides a valid approximation for a matter-dominated universe at early times, $b = a^2$, $\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\langle\rho\rangle$, and $\frac{\ddot{a}}{a} = \frac{4\pi G}{3}\langle\rho\rangle$. Working in comoving coordinates, $\mathbf{x} = \mathbf{r}/a$, one obtains

$$\frac{\dot{x}}{\ddot{x}} = \frac{\dot{a}}{\ddot{a}} = \sqrt{\frac{3}{2\pi G\langle\rho\rangle}}$$

analogously to equation (7), although in this case $\langle\rho\rangle$ depends on time due to the expansion of space (see [14] for a detailed discussion of the cosmological implications of the parallelism between peculiar velocities and accelerations).

The non-local generalization (9) has also received considerable attention during the past years (e.g. [19, 7, 13]). In particular, D. Li, J. Rodrigo and X. Zhang [23] have established local existence (using the linear semigroup and Duhamel's formula), as well as a continuation criterion derived by contradiction (see our Proposition 1) and some results concerning blow-up in finite time (Proposition 3). Global existence when the initial data is small in L^1 has been recently addressed in [25] and [26]. We will show that, in such a case, $\|\rho\|_{L^\infty}(t)$ may be bounded uniformly, and global existence is obtained as a corollary. We state these results as a condition on the size of the pressure forces, β; our sufficient conditions for gravitational stability are related to (and arguably supersede) the classical Jeans criterion, $\beta > 1$ [21], widely applied by the astrophysical comunity.

The paper is organized as follows: In Section 2, we show that the problem is locally well posed (forward in time) in Sobolev spaces (Theorem 1) and a *Beale-Kato-Majda*-type criterion involving $\|\rho\|_{L^\infty}$ (Proposition 1). These results have been proved previously (see Theorems 1.1 and 1.4 in [23]) by different techniques. We also present a result concerning the smoothing effect of our model (Theorem 2), due to the linear part when $\beta > 0$, and prove that the problem is ill posed backward in time. The smoothing effect result, up to our knowledge, is new but it is something that could be expected. The ill-posedness result is almost a consequence of the instant analyticity property. The conditions for gravitational stability, i.e. the decay of small perturbations (Theorem 3) and fluctuations of arbitrary size (Theorem 4), are investigated in Section 3. These results are compared with the well-known critical mass phenomenon in higher dimensions. Using Proposition 1 these theorems implies global in time existence results which are, up to our knowledge, new for equation (9) (compare with Theorem 1.1 in [18]). A finite time blow up result for some initial data in a special class is proved in Theorem 1.10 of [23]. These initial data are concentrated near the origin in terms of a parameter δ such that, for $\delta << 1$, the maximum of the initial datum grows, roughly speaking, like $1/\delta$. Taking this Theorem into account, our global in time result for small data seems to be quite sharp. In this Section we also give a minimum time for blow up which is the periodic version of Theorem 1.9 in [23] . Numerical simulations are presented in Section 4.

2 Well-posedness

2.1 Well-posedness in Sobolev spaces

For the sake of mathematical consistency, we show in this section that equation (9) is well posed if $\beta \geq 0$ and $0 < \alpha \leq 2$. First, let us note that the average density $\langle\rho\rangle$ defined in equation (5) is conserved during the evolution of the system, and thus it is a constant that only depends on the initial data[1]. We state this claim in the following Lemma:

Lemma 1 (Mass conservation). *Let $\rho(x,t)$ be a classical solution. Then, the total mass is conserved*

$$\int_{\mathbb{T}^d} \rho(x,t)\ dx = \int_{\mathbb{T}^d} \rho(x,0)\ dx,$$

and, as a consequence, $\langle\rho\rangle$ is a constant.

Proof. We observe that, using $2\pi\hat{f}(0) = \int_{\mathbb{T}} f(x)\,dx$, the term

$$\int_{\mathbb{T}} \Lambda^\alpha u = 0.$$

Now, integrating and using the Divergence Theorem the proof follows. ☐

To prove the existence and uniqueness of classical solution, we proceed as in [3]. We obtain some *'a priori'* (*i.e.* assuming the existence of classical solution) bounds for the usual norm in the space $H^k(\mathbb{T}^d)$ and then regularize equation (9) so that all the regularized problems have a classical solution for a uniform time T. To conclude, we use the *'a priori'* bound to show that the solutions to the regularized problem form a Cauchy sequence whose limit is the solution to the original equation. In order to simplify the notation, we will abbreviate $\rho(x,t) = \rho(x)$, or simply ρ, throughout the rest of the paper.

---

[1] Without loss of generality, one can set $\langle\rho\rangle = 1$ by an adequate choice of units.

Theorem 1 (Well-posedness). *Given equation (9) with $\beta \geq 0$, $0 < \alpha \leq 2$, and initial data $0 \leq \rho_0(x,0) \in H^k(\mathbb{T}^d)$, where $k > d/2+2$, there exists a time $\tau = \tau(\rho_0) > 0$ and a unique solution $\rho \in C([0,\tau], H^s) \cap L^\infty([0,\tau], H^k(\mathbb{T}))$, for $s < k$. Moreover, $\rho \in C^1([0,\tau], C(\mathbb{T})) \cap C([0,\tau], C^2(\mathbb{T}))$.*

Proof. For simplicity, we only work out in detail the case $k=3$ in one spatial dimension, the general case with $d=2,3$ and $k>3$ being analogous.

(Existence:) We define the norm in $H^k(\mathbb{T})$ as

$$\|\cdot\|_{H^k}^2 = \|\cdot\|_{L^2}^2 + \|\partial_x^k \cdot\|_{L^2}^2.$$

We multiply the equation by ρ and integrate by parts:

$$\int_{\mathbb{T}} \rho(x)\partial_t\rho(x)dx = -\beta\int_{\mathbb{T}} \rho(x)\Lambda^\alpha\rho(x) + \int_{\mathbb{T}}(\rho(x))^2(\rho(x)-1)dx + \int_{\mathbb{T}} \frac{\partial_x(\rho(x))^2}{2}\partial_x U dx$$

Using Parseval Theorem and Hölder inequality we get the estimate

$$\frac{d}{dt}\|\rho\|_{L^2}^2 \leq -2\beta\|\Lambda^{\alpha/2}\rho\|_{L^2}^2 + c_1\|\rho\|_{L^\infty}\|\rho\|_{L^2}^2 \leq c(\|\rho\|_{H^3}+1)^2\|\rho\|_{H^3}. \tag{10}$$

In the same way, for $\|\partial_x^3\rho\|_{L^2}^2$, we obtain the following bound:

$$\frac{d}{dt}\|\partial_x^3\rho\|_{L^2}^2 \leq c(\|\rho\|_{L^\infty} + \|\partial_x^2\rho\|_{L^\infty})\|\rho\|_{H^3}^2 \leq c\|\rho\|_{H^3}(\|\rho\|_{H^3}+1)^2. \tag{11}$$

Using (10) and (11), one finally arrives to

$$\frac{d}{dt}\|\rho\|_{H^3} \leq c(\|\rho\|_{H^3}+1)^2,$$

so

$$\|\rho\|_{H^3} \leq \frac{\|\rho_0\|_{H^3}}{1 - c\|\rho_0\|_{H^3}t}. \tag{12}$$

We only sketch the rest of the proof because it is classical. In order to show the existence of classical solutions to problem (9), we consider $\mathcal{J} \in C_c^\infty$, $\mathcal{J}(x) = \mathcal{J}(|x|)$, $\mathcal{J} \geq 0$, and $\int_{\mathbb{R}} \mathcal{J} = 1$. For $\epsilon > 0$, we define

$$\mathcal{J}_\epsilon(x) = \frac{1}{\epsilon}\mathcal{J}\left(\frac{x}{\epsilon}\right) \tag{13}$$

and consider the regularized problems

$$\partial_t\rho^\epsilon = -\beta\mathcal{J}_\epsilon * \Lambda^\alpha\mathcal{J}_\epsilon * \rho^\epsilon + \mathcal{J}_\epsilon * (\mathcal{J}_\epsilon * \rho^\epsilon(\mathcal{J}_\epsilon * \rho^\epsilon - 1)) + \mathcal{J}_\epsilon * (\nabla\mathcal{J}_\epsilon * \rho^\epsilon \cdot \nabla T(\rho^\epsilon)),$$

where

$$\Delta T(\rho^\epsilon) = (\mathcal{J}_\epsilon * \rho^\epsilon - 1).$$

In these regularized problems, the total mass is also conserved, and the previous bounds (10) and (11) hold. Using classical energy methods (see [3]), one may prove that the regularized problems have an unique smooth solution and, using expression (12), that the sequence ρ^ϵ, indexed in ϵ, is Cauchy in $C([0,\tau], H^s)$, $0 \leq s < 3$, with $\tau = \tau(\rho_0) \leq \frac{1}{c\|\rho_0\|_{H^3}}$. Thus, it has a limit ρ, which is a classical solution of our problem. In order to show that $\rho \in H^3$ we use that ρ^ϵ is uniformly bounded in H^3 and, due to the strong convergence to ρ in H^s with $s < 3$, this implies that ρ^ϵ are weakly convergent to ρ in H^3.

(Uniqueness:) Suppose that for the same initial data ρ_0 we have two classical solutions $\rho_1, \rho_2 \in C([0,\tau], H^3(\mathbb{T}^d))$. Then the equation for $\rho = \rho_2 - \rho_1$ is

$$\partial_t \rho = -\beta\Lambda^\alpha \rho + (\rho_2(\rho_2 - 1) - \rho_1(\rho_1 - 1)) + \nabla\rho_2 \cdot \nabla T(\rho_2) - \nabla\rho_1 \cdot \nabla T(\rho_1),$$

so

$$\frac{1}{2}\frac{d}{dt}\|\rho\|_{L^2}^2 \leq -\beta\|\Lambda^{\alpha/2}\rho\|_{L^2}^2 + \|\rho\|_{L^2}^2(\|\rho_2\|_{L^\infty} + \|\rho_1\|_{L^\infty}) + \int_{\mathbb{T}^d} \rho\nabla\rho_1\nabla(T(\rho_2) - T(\rho_1))dx.$$

We observe that the equation for $T(\rho_2) - T(\rho_1)$ is

$$\Delta(T(\rho_2) - T(\rho_1)) = \rho.$$

Using Poincaré inequality we have that

$$\|\nabla(T(\rho_2) - T(\rho_1))\|_{L^2} \leq c\|\rho\|_{L^2}.$$

Then

$$\int_{\mathbb{T}^d} \rho\nabla\rho_1\nabla(T(\rho_2) - T(\rho_1))dx \leq \|\nabla\rho_1\|_{L^\infty}\|\rho\|_{L^2}\|\nabla(T(\rho_2) - T(\rho_1))\|_{L^2} \leq c\|\nabla\rho_1\|_{L^\infty}\|\rho\|_{L^2}^2.$$

Putting all together we obtain

$$\frac{d}{dt}\|\rho\|_{L^2}^2 \leq c(\rho_2, \rho_1)\|\rho\|_{L^2}^2.$$

Applying Gronwall inequality we conclude the uniqueness. ☐

We give an alternative proof of a *Beale-Kato-Majda*-type criterion characterising the possible singularities (see [23]):

Proposition 1 (Singularity formation)**.** *Suppose that we have*

$$\int_0^{t^+} \|\rho\|_{L^\infty} < \infty,$$

Then the solution exists upon time $t^+ + \sigma$ *for a sufficiently small* σ*.*

Proof. To clarify the exposition, we consider the case with $d = 1$ first. To prove the result we will assume that

$$\int_0^{t^+} \|\rho\|_{L^\infty}(s)ds = M < \infty \tag{14}$$

Using (10) and (11) we have

$$\frac{d}{dt}\|\rho\|_{H^3} \leq c(\|\rho\|_{L^\infty} + \|\partial_x^2\rho\|_{L^\infty} + 1)\|\rho\|_{H^3},$$

so, using Gronwall inequality we conclude that

$$\|\rho\|_{H^3}(t) \leq \|\rho_0\|_{H^3} e^{c\int_0^t \|\rho\|_{L^\infty}(s) + \|\partial_x^2\rho\|_{L^\infty}(s)ds + t}.$$

Now we use the technique exposed in [17]: we use Rademacher Theorem for $\partial_x^2\rho(x_t) = \|\partial_x^2\rho\|_{L^\infty}(t)$. We can do that if the solution is sufficiently smooth. The evolution of this quantity is given by

$$\partial_t\partial_x^2\rho(x_t) = -\beta\Lambda^\alpha\partial_x^2\rho(x_t) + \partial_x^2\rho(x_t)(3(\rho(x_t) - 1) + \rho(x_t)) + 3(\partial_x\rho(x_t))^2.$$

We use the generalized Landau inequality (see [24])

$$|\partial_x\rho(x)|^2 \le 2\rho(x)\|\partial_x^2\rho\|_{L^\infty}.$$

Due to this pointwise inequality we observe that the quadratic terms $(\partial_x\rho(x_t))^2$ are linear in $\partial_x^2\rho(x_t)$. Thus we obtain

$$\partial_t\partial_x^2\rho(x_t) \le 10\|\rho\|_{L^\infty}(t)\partial_x^2\rho(x_t) \Rightarrow \|\partial_x^2\rho\|_{L^\infty}(t) \le \|\partial_x^2\rho_0\|_{L^\infty}e^{10\int_0^t 10\|\rho\|_{L^\infty}(r)dr},$$

and we conclude the result in the one dimensional case .

Now we consider the case $d=2$. The first step is to obtain a bound on $\|\Delta\rho\|_{L^2}$. We multiply the equation by $\Delta^2\rho$ and integrate in space. The dissipative terms are negative so we neglect it. The reaction term contributes with

$$\int_{\mathbb{T}^2}\Delta\rho(x)\Delta(\rho(x)(\rho(x)-1))dx \le c\|\Delta\rho\|_{L^2}^2\|\rho\|_{L^\infty} + \|\Delta\rho\|_{L^2}\||\nabla\rho|^2\|_{L^2},$$

and using a classical Gagliardo-Niremberg interpolation inequality we get

$$\||\nabla\rho|^2\|_{L^2} \le c\|\nabla\rho\|_{L^4}^2 \le c\|\Delta\rho\|_{L^2}\|\rho-1\|_{L^\infty}.$$

Thus the contribution of the reaction term can be bounded as

$$\int_{\mathbb{T}^2}\Delta\rho(x)\Delta(\rho(x)(\rho(x)-1))dx \le c\|\Delta\rho\|_{L^2}^2\|\rho\|_{L^\infty}. \tag{15}$$

The contribution of the transport term is

$$\int_{\mathbb{T}^2}\Delta^2\rho(x)\nabla\rho(x)\cdot\nabla U(x)dx = J_1+J_2+J_3+J_4+J_5,$$

where

$$J_1 = \int_{\mathbb{T}^2}\Delta\rho(x)\nabla U(x)\cdot\nabla\Delta\rho(x)dx \le c\|\Delta\rho\|_{L^2}^2\|\rho\|_{L^\infty},$$

$$J_2 = \int_{\mathbb{T}^2}\Delta\rho(x)|\nabla\rho(x)|^2dx \le c\|\Delta\rho\|_{L^2}\||\nabla\rho|^2\|_{L^2} \le c\|\Delta\rho\|_{L^2}^2\|\rho\|_{L^\infty},$$

$$J_3 = 4\int_{\mathbb{T}^2}\Delta\rho(x)\partial_{x_2}^2\rho(x)\partial_{x_1}^2U(x)dx \le c\|\Delta\rho\|_{L^2}^2\|\partial_{x_1}^2U\|_{L^\infty},$$

$$J_4 = 2\int_{\mathbb{T}^2}(\Delta\rho(x))^2\partial_{x_1}^2U(x)dx \le c\|\Delta\rho\|_{L^2}^2\|\partial_{x_1}^2U\|_{L^\infty},$$

$$J_5 = 2\int_{\mathbb{T}^2}\Delta\rho(x)\partial_{x_2}^2\rho(x)(\rho(x)-1)dx \le c\|\Delta\rho\|_{L^2}^2\|\rho\|_{L^\infty},$$

and, putting all together we obtain

$$\frac{d}{dt}\|\Delta\rho\|_{L^2} \le c\|\Delta\rho\|_{L^2}(\|\rho\|_{L^\infty}+\|\partial_{x_1}^2U\|_{L^\infty}).$$

Due to Gronwall inequality we obtain

$$\|\rho\|_{H^2}(t) \le c(M)\|\rho_0\|_{H^2}\exp\left(\int_0^t\|\partial_{x_1}^2U\|_{L^\infty}(s)+\|\partial_{x_1}\partial_{x_2}U\|_{L^\infty}(s)ds\right). \tag{16}$$

We use a classical inequality (see 3.2b and Appendix 1 in [28] and [2])

$$\|V\|_{L^\infty} \le c(\|\zeta\|_{L^p}+\|\zeta\|_{L^\infty}\max\{1,\log(\|\zeta\|_{W^{s-1,p}}/\|\zeta\|_{L^\infty})\}, \quad 1<p<\infty, s>1+2/p$$

where $V(\zeta)$ is a singular integral operator of Calderon-Zygmund type of ζ. In our case we have

$$\begin{aligned}\|\partial_{x_i}\partial_{x_j}U\|_{L^\infty} &\leq c(\|\rho-1\|_{L^2}+\|\rho-1\|_{L^\infty}\max\{1,\log(\|\rho-1\|_{H^2}/\|\rho-1\|_{L^\infty})\}) \\ &\leq c(M,\|\rho_0\|_{L^2})(1+\|\rho\|_{L^\infty}\max\{1,\log(c\|\rho\|_{H^2})\}). \end{aligned} \tag{17}$$

Inserting in (16) we have

$$\|\rho\|_{H^2}(t)\leq c(M)\|\rho_0\|_{H^2}\exp\left(c(M,\|\rho_0\|_{L^2})\int_0^t(1+\|\rho\|_{L^\infty}(s)\log(c\|\rho\|_{H^2}(s)+e)ds\right).$$

Now let $y(t)=c\|\rho\|_{H^2}(t)+e$. We have

$$y(t)\leq c(M)y(0)\exp\left(c(M,\|\rho_0\|_{L^2})\int_0^t(1+\|\rho\|_{L^\infty}(s)\log(y(s))ds\right),$$

and write $z(t)=\log(y(t))$. We get

$$z(t)\leq c(M)+z(0)+c(M,\|\rho_0\|_{L^2})t+\int_0^t\|\rho\|_{L^\infty}(s)z(s)ds,$$

and due to the integral version of the Gronwall inequality we get

$$z(t)\leq c(M,\|\rho_0\|_{H^2})(1+t). \tag{18}$$

Thus, we conclude that the bound (14) implies $\rho(t)\in H^2$. To obtain higher regularity we can do the same. Indeed, taking derivatives and multiplying by the correct terms we obtain

$$\|\rho\|_{H^4}\leq\|\rho_0\|_{H^4}\exp\left(\int_0^t\|\rho\|_{L^\infty}(s)+\|\Delta\rho\|_{L^2}(s)+\|\partial_{x_1}^2U\|_{L^\infty}(s)+\|\partial_{x_1}\partial_{x_2}U\|_{L^\infty}(s)\right)ds,$$

where we used the following classical interpolation inequalities

$$\|Df\|_{L^4}^2\leq c\|D^2f\|_{L^2}\|f\|_{L^\infty},\ \text{and}\ \|D^2f\|_{L^4}^2\leq\|D^2f\|_{L^2}\|D^2f\|_{L^\infty}.$$

Using the previous bounds (14) (17) and (18) we conclude the result.

□

2.2 Smoothing effect

Our proof follows the steps presented in [16]. The proof is based in *a priori* estimates for the complex extension of the function ρ on the complex strip $S=\{x+i\xi,x\in\mathbb{T},|\xi|<kt\}$, for certain constant $k>0$ (see Figure 1). We define

$$\|\rho\|_{L^2(S)}^2=\int_{\mathbb{T}}|\rho(x+ikt)|^2dx+\int_{\mathbb{T}}|\rho(x-ikt)|^2dx,\quad \|\rho\|_{H^3(S)}^2=\|\rho\|_{L^2(S)}^2+\|\partial_x^3\rho\|_{L^2(S)}^2.$$

For the complex extension the equation is

$$\partial_t\rho(x+i\xi)=-\beta\Lambda^\alpha\rho(x+i\xi)+\rho(x+i\xi)(\rho(x+i\xi)-1)+\nabla\rho(x+i\xi)\cdot\nabla T(\rho)(x+i\xi) \tag{19}$$

Theorem 2 (Smoothing effect)**.** *Let us consider equation (9) with $\alpha\geq1$ with $\rho_0\in H^3$ as initial data and $\mathbb{T}$ as spatial domain. Then the classical solution ρ (which exists, at least locally in time, because of Theorem 1) continues analitically into the strip S for times $0<t\leq\tau(\rho_0)$.*



Figure 1: The strip of analyticity for ρ.

Proof. For the sake of brevity we work with both terms $\pm ikt$ at the same time. Firstly we consider the critical case $\alpha = 1$. We start with the $L^2(S)$ norm:

$$\frac{1}{2}\frac{d}{dt}\int_{\mathbb{T}} |\rho(x \pm ikt)|^2 dx = \text{Re}\int_{\mathbb{T}} \bar{\rho}(x \pm ikt)(\partial_t \rho(x \pm ikt) \pm ik\partial_x \rho(x \pm ikt))dx.$$

Using Plancherel Theorem we have

$$-\beta \int_{\mathbb{T}} \bar{\rho}(x \pm ikt)\Lambda\rho(x \pm ikt)dx = -\beta \int_{\mathbb{T}} |\Lambda^{1/2}\rho(x \pm ikt)|^2 dx \leq 0.$$

The reaction term can be bounded as follows:

$$\text{Re}\int_{\mathbb{T}} \bar{\rho}(x \pm ikt)\rho(x \pm ikt)(\rho(x \pm ikt) - 1)dx \leq (\|\rho\|_{L^\infty(S)} + 1)\|\rho\|^2_{L^2(S)}.$$

We have

$$\begin{aligned} I_1 = \text{Re}\int_{\mathbb{T}} \bar{\rho}(x \pm ikt)\partial_x\rho(x \pm ikt)\partial_x U(x \pm ikt)dx &= \text{Re}\int_{\mathbb{T}} \partial_x U(x \pm ikt)\frac{1}{2}\partial_x|\rho(x \pm ikt)|^2 dx \\ &+ \text{Re}\, i\int_{\mathbb{T}} \partial_x U(x \pm ikt)\text{Re}\rho(x \pm ikt)\partial_x \text{Im}\rho(x \pm ikt)dx \\ &- \text{Re}\, i\int_{\mathbb{T}} \partial_x U(x \pm ikt)\text{Im}\rho(x \pm ikt)\partial_x \text{Re}\rho(x \pm ikt)dx. \end{aligned}$$

Integrating by parts we have

$$\begin{aligned} I_1 = \text{Re}\int_{\mathbb{T}} &\partial_x U(x \pm ikt)\frac{1}{2}\partial_x|\rho(x \pm ikt)|^2 dx \\ &- \text{Re}\, i\int_{\mathbb{T}} \partial_x^2 U(x \pm ikt)\text{Re}\rho(x \pm ikt)\text{Im}\rho(x \pm ikt)dx \\ &+ 2\int_{\mathbb{T}} \partial_x \text{Im}U(x \pm ikt)\text{Im}\rho(x \pm ikt)\partial_x \text{Re}\rho(x \pm ikt)dx, \end{aligned}$$

and we have

$$I_1 \leq c\|\rho\|^2_{L^2(S)}(\|\rho\|_{L^\infty(S)}+1)+\|\partial_x \mathrm{Im}U\|_{L^\infty(S)}\|\rho\|_{L^2(S)}\|\partial_x\rho\|_{L^2(S)}.$$

Due to the mean conservation and that the initial data has $\mathrm{Im}\rho_0=0$ we have the bound

$$\|\partial_x \mathrm{Im}U\|_{L^\infty(S)} \leq c\|\partial_x^2 \mathrm{Im}U\|^{1/2}_{L^2(S)}\|\partial_x \mathrm{Im}U\|^{1/2}_{L^2(S)} \leq c\|\mathrm{Im}\rho\|_{L^2(S)},$$

where in the last inequality we use the Poincaré inequality.

The last term can be bounded as follows:

$$\mathrm{Re}\int_{\mathbb{T}} \bar\rho(x\pm ikt)(\pm ik\partial_x\rho(x\pm ikt))dx \leq k\|\rho\|_{L^2(S)}\|\partial_x\rho\|_{L^2(S)}.$$

Putting all together and using Sobolev embedding we have

$$\frac{d}{dt}\|\rho\|^2_{L^2(S)} \leq c\|\rho\|^2_{H^3(S)}(1+\|\rho\|_{H^3(S)}). \tag{20}$$

We have

$$\frac{1}{2}\frac{d}{dt}\int_{\mathbb{T}}|\partial_x^3\rho(x\pm ikt)|^2dx = \mathrm{Re}\int_{\mathbb{T}}\partial_x^3\bar\rho(x\pm ikt)(\partial_t\partial_x^3\rho(x\pm ikt)\pm ik\partial_x^4\rho(x\pm ikt))dx.$$

Following the previous techniques we have

$$\begin{aligned}\frac{1}{2}\frac{d}{dt}\int_{\mathbb{T}}|\partial_x^3\rho(x\pm ikt)|dx \leq c\|\rho\|^2_{H^3(S)}(\|\rho\|_{H^3(S)}+1)-\beta\int_{\mathbb{T}}|\Lambda^{1/2}\partial_x^3\rho|^2dx\\ +\mathrm{Re}\int_{\mathbb{T}}\partial_x^3\bar\rho\partial_x^4\rho\partial_x U dx \pm k\mathrm{Re}\, i\int_{\mathbb{T}}\partial_x^3\bar\rho\partial_x^4\rho dx.\end{aligned}$$

We write

$$J_1=\mathrm{Re}\int_{\mathbb{T}}\partial_x^3\bar\rho\partial_x^4\rho\partial_x U dx \quad \text{and } J_2=\pm k\mathrm{Re}\, i\int_{\mathbb{T}}\partial_x^3\bar\rho\partial_x^4\rho dx.$$

Splitting into real and imaginary part we have

$$\begin{aligned}J_1=\mathrm{Re}\int_{\mathbb{T}}\partial_x U\frac{1}{2}\partial_x|\partial_x^3\rho|^2dx+\int_{\mathbb{T}}\partial_x^3\mathrm{Re}\rho\partial_x^4\mathrm{Im}\rho\mathrm{Im}\partial_x U dx-\int_{\mathbb{T}}\partial_x^3\mathrm{Im}\rho\partial_x^4\mathrm{Re}\rho\mathrm{Im}\partial_x U dx\\ \leq c\|\partial_x^3\rho\|_{L^2(S)}(\|\rho\|_{L^\infty(S)}+1)+2K_1,\end{aligned}$$

where the last integral is

$$\begin{aligned}K_1&=\left|\int_{\mathbb{T}}\partial_x^3\mathrm{Re}\rho\Lambda H\partial_x^3\mathrm{Im}\rho\mathrm{Im}\partial_x U dx\right|\\ &=\left|\int_{\mathbb{T}}\Lambda^{1/2}H\partial_x^3\mathrm{Im}\rho\Lambda^{1/2}(\partial_x^3\mathrm{Re}\rho\mathrm{Im}\partial_x U)dx\right|\\ &\leq \|\Lambda^{1/2}\partial_x^3\rho\|_{L^2(S)}\|\Lambda^{1/2}(\partial_x^3\mathrm{Re}\rho\mathrm{Im}\partial_x U)\|_{L^2(S)}.\end{aligned}$$

We use the commutator estimate

$$\|\Lambda^{1/2}(fg)-f\Lambda^{1/2}g\|_{L^2(S)} \leq c\|\partial_x f\|_{L^\infty(S)}\|g\|_{L^2(S)}$$

to conclude that

$$\begin{aligned}K_1&\leq \|\Lambda^{1/2}\partial_x^3\rho\|_{L^2(S)}(\|\partial_x^2\mathrm{Im}U\|_{L^\infty(S)}\|\partial_x^3\rho\|_{L^2(S)}+\|\partial_x\mathrm{Im}U\|_{L^\infty(S)}\|\Lambda^{1/2}\partial_x^3\rho\|_{L^2(S)})\\ &\leq c(\|\Lambda^{1/2}\partial_x^3\rho\|_{L^2(S)}\|\rho\|_{L^\infty(S)}\|\partial_x^3\rho\|_{L^2(S)}+\|\mathrm{Im}\rho\|_{L^2(S)}\|\Lambda^{1/2}\partial_x^3\rho\|^2_{L^2(S)}).\end{aligned}$$

If we use Young inequality we have

$$K_1 \leq \left(\frac{\epsilon}{2} + c\|\mathrm{Im}\rho\|_{L^2(S)}\right) \|\Lambda^{1/2}\partial_x^3\rho\|^2_{L^2(S)} + \frac{1}{2\epsilon}\|\rho\|^2_{L^\infty(S)}\|\partial_x^3\rho\|^2_{L^2(S)}.$$

We can also bound

$$J_2 \leq k\|\Lambda^{1/2}\partial_x^3\rho\|_{L^2(S)}.$$

Finally, taking $\epsilon = \beta/2$

$$\begin{aligned}\frac{d}{dt}\int_{\mathbb{T}} |\partial_x^3\rho(x \pm ikt)|dx \leq c\|\rho\|^2_{H^3(S)}(\|\rho\|_{H^3(S)} + 1)\\ + \left(c\|\mathrm{Im}\rho\|_{L^2(S)} + k - \frac{3\beta}{4}\right) \|\Lambda^{1/2}\partial_x^3\rho\|_{L^2(S)}, \quad (21)\end{aligned}$$

and this is a correct bound if the term

$$K_2(t) = c\|\mathrm{Im}\rho\|_{L^2(S)} + k - \frac{3\beta}{4} \leq 0.$$

Initially $K_2(0) = k - \frac{3\beta}{4}$, so taking $k = \beta/4 < \beta$ the condition $K_2 < 0$ will be satisfied for a (maybe short) time if we can bound the evolution of $\|\mathrm{Im}\rho\|_{L^2(S)}$.

We define the *'energy'*

$$\|\rho\|_{A(S)} = \|\rho\|_{H^3(S)} + \frac{1}{\frac{\beta}{2} - c\|\mathrm{Im}\rho\|_{L^2(S)}}.$$

The evolution of this energy is

$$\frac{d}{dt}\|\rho\|_{A(S)} \leq c((\|\rho\|_{A(S)} + 1)^2 + \|\rho\|^2_{A(S)}\frac{d}{dt}\|\mathrm{Im}\rho\|_{L^2(S)}).$$

At this point it is easy to obtain

$$\frac{d}{dt}\|\mathrm{Im}\rho\|_{L^2(S)} \leq c(\|\rho\|_{H^3(S)} + 1)\|\mathrm{Im}\rho\|^2_{L^2(S)},$$

so

$$\frac{d}{dt}\|\rho\|_{A(S)} \leq (\|\rho\|_{A(S)} + 1)^5$$

and we conclude that there is a time $\tau(\rho)$ so that $\|\rho\|_{A(S)} < \infty$.

Now we approximate the problem in the real line as in the proof of the Theorem 1, using the rescaled heat kernel as a mollifier:

$$\partial_t\rho^\epsilon = -\beta\mathcal{J}_\epsilon * \Lambda^\alpha\mathcal{J}_\epsilon * \rho^\epsilon + \mathcal{J}_\epsilon * (\mathcal{J}_\epsilon * \rho^\epsilon(\mathcal{J}_\epsilon * \rho^\epsilon - 1)) + \mathcal{J}_\epsilon * (\nabla\mathcal{J}_\epsilon * \rho^\epsilon \cdot \nabla\mathcal{J}_\epsilon * T(\rho^\epsilon)),$$

where

$$\Delta T(\rho^\epsilon) = (\mathcal{J}_\epsilon * \rho^\epsilon - 1),$$

and we use $\rho_0^\epsilon = \mathcal{J}_\epsilon * \rho_0$ as initial data. Using Picard's Theorem in $H^3(\mathbb{T})$ we obtain solutions ρ^ϵ up to time τ^ϵ which are analytic. Due to the above estimates we have a time a common time of existence τ depending on the initial data ρ_0 but not in ϵ and if $t < \tau(\rho_0)$, $\|\rho^\epsilon\|_{H^3(S)} < \infty$. Now we pass to the limit in the Hardy-Sobolev space $H^3(S)$ obtaining ρ and we use the fact that if the energy $\|\cdot\|_{A(S)}$ remains bounded then this implies the analyticity. Due to the uniqueness, this is the same ρ as in Theorem 1.

In the proof for the case $\alpha > 1$ we use the inequality

$$\|\Lambda^{1/2}T\|^2_{L^2(S)} \leq \|T\|^2_{L^2(S)} + \|\Lambda^{\alpha/2}T\|^2_{L^2(S)}$$

obtained by splitting in the Fourier side. We conclude the proof in the supercritical case following the same steps. □

Using the smoothing effect, one can prove ill-posedness in Sobolev spaces when the problem is considered backward in time. We remark that this result does not require global existence of solutions.

Proposition 2 (Ill-posedness)**.** *There are solutions $\tilde{\rho}$ to the backward in time equation* (9) *with* $\beta > 0$*, such that* $\|\tilde{\rho}\|_{H^k}(0) < \epsilon$ *and* $\|\tilde{\rho}\|_{H^k}(\delta) = \infty$ *for all* $0 < \epsilon$*, sufficiently small* $0 < \delta$*, and* $k > 3$*.*

Proof. Let us fix $k = 4$, the other cases being similar. Take $g_0(x) \in H^3(\mathbb{T})$ but $g_0 \notin H^4(\mathbb{T})$ and consider the solution (forward in time) ρ^λ to the equation (9) with initial data $\rho(x,0) = \lambda g_0(x)$ where $0 < \lambda < 1$. We have (see Theorem 2) that there is a uniform (in λ) time $\delta^*(g_0)$ such that ρ^λ exists and is analytic up to time $\delta^*(g_0)$. Now define $\tilde{\rho}^{\lambda,\delta}(x,t) = \rho^\lambda(x, -t+\delta)$ for fixed $0 < \delta < \delta^*(g_0)$. We have that $\|\tilde{\rho}^{\lambda,\delta}\|_{H^4}(\delta) = \lambda\|g_0\|_{H^4} = \infty$. Now, using Theorem 2 we have that there is a growing strip of analitycity for ρ^λ where we can apply Cauchy's integral formula in order to obtain

$$\|\partial_x^4 \tilde{\rho}^{\lambda,\delta}\|_{L^2(\mathbb{T})}(0) = \|\partial_x^4 \rho^\lambda\|_{L^2(\mathbb{T})}(\delta) \leq \frac{C(\beta)}{\delta^*(g_0)}\|\partial_x^3 \rho^\lambda\|_{L^2(S_{\delta^*})} \leq \frac{c(\beta, g_0)\lambda}{\delta}.$$

Taking $0 < \lambda < \min\{1, \epsilon\delta/c(\beta, g_0)\}$ we conclude the proof. ☐

3 Gravitational stability

The blow-up of the solutions in finite time is of particular interest from the physical point of view, since it determines whether a gravitationally bound structure may form or not. In order to investigate the necessary conditions for star formation, the present section is devoted to the evolution of $\|\rho\|_{L^\infty}(t)$. More precisely, it is our goal to determine whether/when the growth of singularities can be avoided in our non-local adhesion model. We will focus on the critical case ($\alpha = 1$) and remark that the resulting equation is more dissipative if $\alpha > 1$. All our results are expressed in units such as $\langle\rho\rangle = 1$; the conversion to other systems can be trivially carried out by substituting every occurrence of ρ by $\frac{\rho}{\langle\rho\rangle}$ or, alternatively, β by $\beta\langle\rho\rangle$.

3.1 Minimum time for star formation

First of all, let us address the minimum time required for the gravitational collapse of a protostar. For those cases where a star is formed, i.e. $\rho(x_*, t_*) \to \infty$, we provide a lower bound for the value of t_* as a function of the initial condition. Note that, by virtue of Proposition 1, classical solution is guaranteed up to a time t_*.

Proposition 3 (Minimum time for blow-up)**.** *Let* $\rho \geq 0$ *be a smooth solution of* (9) *in* $\mathbb{T}^d$, $d \leq 3$*. Then we have*

$$\frac{d}{dt}\|\rho\|_{L^\infty}(t) \leq \|\rho\|_{L^\infty}(t)(\|\rho\|_{L^\infty}(t) - 1),$$

thus

$$\|\rho\|_{L^\infty}(t) \leq \frac{\|\rho_0\|_{L^\infty}}{\|\rho_0\|_{L^\infty} + (1 - \|\rho_0\|_{L^\infty})e^t}.$$

As a consequence the minimum time for the formation of a star is

$$t^* = \log\left(\frac{\|\rho_0\|_{L^\infty}}{\|\rho_0\|_{L^\infty} - 1}\right).$$

Proof. Due to the smoothness of ρ and using Rademacher Theorem, we have that the evolution of

$$\|\rho\|_{L^\infty}(t) = \max_x \rho(x,t) = \rho(x_t),$$

is given by

$$\frac{d}{dt}\rho(x_t) \leq -\beta\Lambda^\alpha \rho(x_t) + \rho(x_t)(\rho(x_t)-1) \leq \rho(x_t)(\rho(x_t)-1),$$

where the last inequality is due to the integral representation for the diffusive operator. Now the result is straightforward. ☐

3.2 Exponential suppresion of density fluctuations

Our main result, and the one that bears more physical relevance, is that equation (9) with $\alpha = 1$ does not have a blow-up for $\|\rho\|_{L^\infty}(t)$ – i.e. no star can ever form – if β is big enough. Physically, β is the dimensionless ratio of the free-fall time, representative of the strength of the gravitational forces, and the sound-crossing time, representative of the restoring pressure force. Depending on the amplitude of the initial conditions (more precisely, the value of $\|\rho_0\|_{L^\infty}$), there is a critical value of the sound speed above which the pressure gradient is able to overcome gravity. Or, in other words, for a uniform gas cloud at a given temperature (i.e. β), sufficiently small perturbations cannot grow. Mathematically, this result is a Maximum Principle for $\|\rho\|_{L^\infty}$ for a special class of initial data, and the description in terms of β or $\|\rho_0\|_{L^1}$ merely reflects the choice of units. Both choices are, of course, absolutely equivalent.

Consistently with the results of numerical simulations, we find that, for sufficiently small perturbations and/or large pressure forces, there is a depletion of the solution (see Figure 2), and the system evolves towards the homogeneous equilibrium state $\rho(x) = 1$. Since $\|\rho\|_{L^\infty}$ is bounded, global existence of classical solution follows from Proposition 1.



Figure 2: a) The initial data (in blue) and the evolution (in red) for the one-dimensional case of equation (9) with $\beta = 1$ and b) the complete evolution.

Theorem 3 (Decay of small perturbations)**.** *For* (9) *with* $0 < \alpha < 2$*, spatial domain* $\mathbb{T}^d$*,* $d = 1, 2, 3$*, and initial data* ρ_0*, if*

$$\|\rho_0\|_{L^\infty} \leq c_{\alpha,d}\beta,$$

where $c_{\alpha,d}$ *is an explicit constant, then the maximum density decays exponentially,*

$$\|\rho\|_{L^\infty}(t) \;\leq\; 1 + \left[\,\|\rho_0\|_{L^\infty} - 1\,\right]\; e^{-\left[c_{\alpha,d}\beta - \|\rho_0\|_{L^\infty}\right]t}.$$



Figure 3: a)The initial data (in blue) and the evolution (in red) for the one-dimensional case of equation (9) with $\beta = 0$ and b) the total evolution.

Moreover, we have that $c_{1,1} = 1$.

Proof. First, we prove the case $\alpha = 1$, $d = 1$. Denoting the point where ρ achieves its maximum value by x_t and using the integral representation of the fractional Laplacian, one can easily derive the bound

$$\Lambda\rho(x_t) = \frac{1}{2\pi}\text{P.V.}\int_{\mathbb{T}} \frac{\rho(x_t)-\rho(y)}{\sin^2\left(\frac{x_t-y}{2}\right)}dy \ \geq \ \frac{1}{2\pi}\int_{\mathbb{T}} \rho(x_t)-\rho(y)\,dy = \rho(x_t)-1$$

Thus

$$\partial_t\rho(x_t) \ \leq \ -\beta\,[\,\rho(x_t)-1\,] + \rho(x_t)\,[\,\rho(x_t)-1\,] \ = \ [\,\rho(x_t)-1\,]\,[\,\rho(x_t)-\beta\,],$$

and the maximum density will always decrease if $\rho(x_t) < \beta$. In fact,

$$\partial_t\,[\,\rho(x_t)-1\,] \ \leq \ -\,[\,\beta - \|\rho_0\|_{L^\infty}\,]\,[\,\rho(x_t)-1\,]\,.$$

Now we proceed with the $d-$dimensional case. From the integral representation of the fractional Laplacian in $\mathbb{T}^d$

$$\Lambda^\alpha\rho(x_t) = C_{\alpha,d}\sum_{\nu\in\mathbb{Z}^d}\text{P.V.}\int_{\mathbb{T}^d}\frac{\rho(x_t)-\rho(y)}{|x_t-y-2\pi\nu|^{d+\alpha}}dy$$

we have that

$$\Lambda^\alpha\rho \geq C_{\alpha,d}\ \text{P.V.}\int_{\mathbb{T}^d}\frac{\rho(x_t)-\rho(y)}{|x_t-y|^{d+\alpha}}dy.$$

Since $|x_t-y| \leq \pi\sqrt{d}$,

$$\Lambda^\alpha\rho \geq c_{\alpha,d}\,[\rho(x_t)-1]\,.$$

and we can conclude the result analogously to the one dimensional case. ☐

In one spatial dimension, it is also possible to prove that, for sufficiently large β (or, equivalently, sufficiently small $\|\rho_0\|_{L^1}$), *all* density fluctuations are exponentially suppressed, regardless of their initial amplitude.

Theorem 4 (Global stability for $d = 1$)**.** *Given equation* (9) *with* $\alpha = 1$*, spatial domain* $\mathbb{T}$*, and initial data* ρ_0*, if*

$$\beta \geq 4\pi^2,$$

then the maximum density decays exponentially,

$$\|\rho\|_{L^\infty}(t) \;<\; 1 + [\,\|\rho_0\|_{L^\infty} - 1\,]\; e^{-t}.$$

Moreover, if $1<\alpha<2$ *then, for any* $\beta>0$*, the following inequality holds*

$$\|\rho\|_{L^\infty} \le c(\alpha,\beta).$$

Proof. Let us consider the neighbourhood around the absolute density maximum

$$\Omega_\delta = \{\, y \mid |x_t - y| \le 2\delta \,\},$$

where δ is a parameter that will be set later, and decompose it in two subsets, $\Omega_\delta = \mathcal{U}_1 \cup \mathcal{U}_2$, such that

$$\mathcal{U}_1 = \{\, y \in \Omega_\delta \mid \rho(x_t) - \rho(y) \ge \rho(x_t)/2 \,\},$$

and

$$\mathcal{U}_2 = \{\, y \in \Omega_\delta \mid \rho(x_t) - \rho(y) < \rho(x_t)/2 \,\}.$$

Since $|x| \ge |\sin(x)|$,

$$\Lambda\rho(x_t) \;\ge\; \frac{1}{2\pi}\int_{\Omega_\delta} \frac{\rho(x_t)-\rho(y)}{\left(\frac{x_t-y}{2}\right)^2}dy \;\ge\; \frac{1}{2\pi}\int_{\mathcal{U}_1} \frac{\rho(x_t)/2}{\delta^2}dy \;=\; \frac{1}{2\pi}\frac{\rho(x_t)}{2\delta^2}\left(|\Omega_\delta| - |\mathcal{U}_2|\right).$$

Substituting $|\Omega_\delta| = 4\delta$ and using that

$$2\pi = \int_{\mathbb{T}} \rho(y)dy \;>\; \frac{\rho(x_t)}{2}|\mathcal{U}_2|,$$

one obtains the lower bound

$$\Lambda\rho(x_t) \;>\; \frac{1}{2\pi}\frac{\rho(x_t)}{2\delta^2}\left[4\delta - \frac{4\pi}{\rho(x_t)}\right],$$

which is maximized for the choice

$$\delta = \frac{2\pi}{\rho(x_t)}.$$

It is important to note, though, that the compactness of the domain also imposes that $\delta \le \pi/2$ or, in other words,

$$\rho(x_t) \ge 4.$$

Inserting the bound for the dissipative term in equation (9), we have

$$\partial_t\rho(x_t) \le -\beta\,\frac{\rho^2(x_t)}{4\pi^2} + \rho(x_t)\,[\,\rho(x_t) - 1\,] = \rho^2(x_t)\left[1 - \frac{\beta}{4\pi^2}\right] - \rho(x_t)$$

If $\beta \ge 4\pi^2$,

$$\partial_t\rho(x_t) \;\le\; -\rho(x_t) \;<\; -[\,\rho(x_t) - 1\,],$$

which proves the desired result for $\rho(x_t) \ge 4$. Theorem 3 implies that this bound (indeed, a stricter one) also applies for $\rho(x_t) < 4$. This concludes with the first part of the result. In the case $1<\alpha<2$ we have

$$\Lambda\rho(x_t) \;>\; \frac{1}{2\pi}\frac{\rho(x_t)}{2\delta^{1+\alpha}}\left[4\delta - \frac{4\pi}{\rho(x_t)}\right].$$

Consequently, we get

$$\partial_t\rho(x_t) \le -\beta\,\frac{\rho^{1+\alpha}(x_t)}{(2\pi)^{1+\alpha}} + \rho(x_t)\,[\,\rho(x_t) - 1\,],$$

and we conclude the proof. □

`Remark 1` Recall that in 2D (8) has a critical mass phenomenon, *i.e.* if $\|\rho_0\|_{L^\infty(\mathbb{R}^2)} < 8\pi$, then there is a global solution, while if $\|\rho_0\|_{L^\infty(\mathbb{R}^2)} > 8\pi$ there is a finite time blow up. If we take $\beta = 1$ and an initial datum with arbitrary L^1 norm, Theorem 4 gives us that, if $\|\rho_0\|_{L^1} \leq \frac{1}{2\pi}$ then the maximum decays exponentially and the solution is global.

4 Numerical simulations

In order to simulate the equation (9) we use a Fourier-collocation method (see [15]). We use Fast Fourier Transform techniques to solve the Poisson equation and to approximate the spatial part of (9). For the time integrator we use an explicit Runge-Kutta of order 4.

We center our atention in the 2D case. The other cases are analogous. We consider an equispaced discretization of our spatial domain $\mathbb{T}^2$,

$$\mathcal{T}_N^2 = \left\{(x_k, y_l) \,:\, x_k = \frac{\pi}{N}(2k - N)\,,\, y_l = \frac{\pi}{N}(2l-1)\,,\, k, l = 0, \dots, N-1\right\},$$

where N is the number of points on each interval $[-\pi, \pi]$. We consider as our approximate solution the pair $\rho(x, y)$ and $T(\rho)(x, y)$ in the grid $\mathcal{T}_N^2$. We have

$$\rho(x_k, y_l) = \sum_{n,m=-N/2}^{N/2-1} \tilde{\rho}_{n,m} e^{in\,x_k} e^{im\,y_l}, \quad -(n^2+m^2)\tilde{T(\rho)}_{n,m} = 4\pi G \tilde{\rho}_{n,m}, \quad \tilde{T(\rho)}_{0,0} = 0,$$

and it is important to remark that the coefficients $\tilde{\rho}_{n,m}$ are related to the so-known Fourier coefficients, typically $\hat{\rho}_{n,m}$, but they are not the same (see [15]). In the same way we can approximate all terms in (8). Once we compute derivatives of ρ and U via the FFT in the equation, we get back to the physical space using the IFFT.

This ends the space discretization part. We consider our equation in the time interval $[0, \tau]$, for some $\tau > 0$. As before, let $\mathcal{G}_J$ an equispaced partition of the time interval,

$$\mathcal{G}_J = \{t_s \,:\, t_s = s\tau/J,\, s = 0, \dots, J-1\},$$

where J is the number points in the time grid. For the evolution part of the equation, $\partial_t \rho$, we are going to use a Runge-Kutta method (specifically, the classic explicit method of order 4), and we start with the initial data

$$\rho_0(x, y) = |\sin x| + |\sin y|$$

represented in Figure 4.

We studied the evolution of $\|\rho\|_{L^\infty}(t)$ for various $\beta \in [0, 1]$. On the left panel in Figure 5, we highlight in red the case $\beta = 0.2$ and in blue $\beta = 0.8$. On the right panel, we plot the results for values of β in the interval $[0.32, 0.34]$.

The mechanism for singularity formation, i.e. the blow-up for $\|\rho\|_{L^\infty}(t)$, is illutrated in Figure 6. In Figure 7, one can see how the solution is depleted for large β.

References

[1] J. M. Bardeen, J. R. Bond, N. Kaiser and A. S. Szalay, *The statistics of peaks of Gaussian random fields*, Astrophysical Journal **304**, 15-61, 1986.

[2] J.T.Beale, T. Kato, and A. Majda, *Remarks on the breakdown of smooth solutions for the 3-D Euler equations*, Communications in Mathematical Physics, **94**, no. 1, 61-66, 1984.



Figure 4: Initial data for ρ and v with $\beta = 10$.



Figure 5: a) The evolution of $\|\rho\|_{L^\infty}(t)$ for $\beta = 0.2$ up to 0.8, with final time $T = 1$, and b) the evolution of $\|\rho\|_{L^\infty}(t)$ for $\beta = 0.32$ up to 0.34, with final time $T = 9$.

[3] A.L. Bertozzi and A.J. Majda, Vorticity and the Mathematical Theory of Incompressible Fluid Flow. *Cambridge Press*, 2002.

[4] P. Biler and T. Nadzieja, *Existence and nonexistence of solutions for a model of gravitational interaction of particles (part 1).*, Colloq. Math. **66**, no. 2, 319–334, 1994.

[5] P. Biler, D. Hilhorst and T. Nadzieja, *Existence and nonexistence of solutions for a model of gravitational interaction of particles (part 2).*, Colloq. Math. **67**, no. 2, 297–308, 1994.

[6] P. Biler, *Existence and nonexistence of solutions for a model of gravitational interaction of particles (part 3).*, Colloq. Math. **68**, no. 2, 229–239, 1995.

[7] P. Biler, G. Karch, *Blowup of solutions to generalized Keller–Segel model*, Journal of Evolution Equations **10**, no. 2, 247-262, 2010.

[8] P. Biler, G. Karch, P. Laurençot and T. Nadzieja, *The 8π-problem for radially symmetric solutions of a chemotaxis model in the plane.*, Math Meth. Appl. Sci. **29**, no. 13, 1563–1583, 2006.

[9] A. Blanchet., *On the parabolic-elliptic Patlak-Keller-Segel system in dimension 2 and higher*, Arxiv preprint arXiv:1109.1543, 2011.



Figure 6: a) ρ for time $t = 0.5$ and $\beta = 0$ and b) v at the same time.



Figure 7: a) ρ for time $t = 1$ and $\beta = 0.5$ and b) v at the same time.

[10] A. Blanchet, J.A. Carrillo and N. Masmoudi, *Infinite time aggregation for the critical Patlak-Keller-Segel model in* $\mathbb{R}^2$, Comm. Pure and Appl. Math. **61**, no. 10, 1449–1481, 2008.

[11] A. Blanchet, J.A. Carrillo. and P. Laurençot, *Critical mass for a Patlak–Keller–Segel model with degenerate diffusion in higher dimensions*, Calculus of Variations and Partial Differential Equations **35**, no. 2, 133-168, 2009.

[12] A. Blanchet, E. Carlen, J.A. Carrillo, *Functional inequalities, thick tails and asymptotics for the critical mass Patlak-Keller-Segel model*, preprint arXiv:1009.0134, 2010.

[13] N. Bournaveas and V. Calvez, *The one-dimensional Keller–Segel model with fractional diffusion of cells*, Nonlinearity, **26**, 923, 2010.

[14] T. Buchert and A. Domínguez, *Adhesive gravitational clustering*, Astron. Astrophys. **438**, 443–460, 2005

[15] Canuto, C., Hussaini, M. Y., Quarteroni, A., Zang, T.A., *Spectral Methods*, Springer-Verlag, 2006

[16] A. Castro, D. Cordoba, C. Fefferman, F. Gancedo, M. López-Fernández, *Rayleigh-Taylor breakdown for the Muskat problem with applications to water waves, To appear in Annals of Mathematics*

[17] A. Córdoba and D. Córdoba, *A maximum principle applied to Quasi-geostrophic equations*, Comm. Math. Phys. **249**, no. 3, 511-528, 2004.

[18] L. Corrias, and B. Perthame and H. Zaag, *Global solutions of some chemotaxis and angiogenesis systems in high space dimensions*, Milan Journal of Mathematics, **72**, no. 1, 1-28, 2004.

[19] C. Escudero, *The fractional Keller-Segel model.* Nonlinearity 19. 2909. 2006.

[20] W. Jäger and S. Luckhaus, *On explosions of solutions to a system of partial differential equations modelling chemotaxis*, Transactions of the American Mathematical Society **329**, 819–824, 1992.

[21] J. H. Jeans, *The Stability of a Spherical Nebula*, Philosophical Transactions of the Royal Society of London. Series A, Containing Papers of a Mathematical or Physical Character **199**, 1–53, 1902.

[22] E. F. Keller and L. A. Segel, *Initiation of slime mold aggregation viewed as an instability*, Journal of Theoretical Biology, **26**, 399–415, 1970.

[23] D. Li, J. Rodrigo and X. Zhang, *Exploding solutions for a nonlocal quadratic evolution problem*, Rev. Mat. Iberoamericana, **26**, 295–332, 2010.

[24] V. Maz'ya, and T. Shaposhnikova, *On pointwise interpolation inequalities for derivatives.* Mathematica Bohemica, **124**, 131-148, 1999.

[25] T. Nagai, *Global existence and decay estimates of solutions to a parabolic-elliptic system of drift-diffusion type in* $\mathbb{R}^2$ Differential Integral Equations **24**, no. 1-2, 2968, 2011.

[26] T. Nagai, *A parabolic-elliptic system of drift-diffusion type with subcritical mass in* $\mathbb{R}^2$. To appear.

[27] C. S. Patlak, *Random walk with persistence and external bias*, Bulletin of Mathematical Biology, **15**, 311–338, 1953.

[28] T. Kato, and G. Ponce, *Well-Posedness of the Euler and Navier-Stokes Equations in the Lebesgue Spaces* $L^p_s(\mathbb{R}2)$, Revista matemática iberoamericana, **2**, no. 1, 73-88, 1986.

[29] , T. Senba, and T. Suzuki, *Weak solutions to a parabolic-elliptic system of chemotaxis*, Journal of Functional Analysis, **191**, no.1, 17-51, 2002.

[30] Ya. B. Zel'Dovich, *Gravitational instability: An approximate theory for large density perturbations.* Astron. Astrophys. **5**, 84–89, 1970.